

Child and Adolescent Mental Health Care in Uganda

This report was developed by ICHAD (International Center for Child Health and Development), SMART Africa Center (Strengthening Mental health And Research Training in Sub-Saharan Africa), the Clark-Fox Policy Institute at the Brown School at Washington University in St. Louis and ChildFund Uganda.

In Uganda, children make up about half (56%) of the total population, and they often present with multiple physical, mental health, and educational challenges.^{1,2} Large numbers of Ugandan children live in communities with high rates of chronic poverty (38%), domestic violence (30%), physical violence toward children (80%), depression (33 to 39%), malaria (70 to 80%), and HIV or AIDS (6%).^{3,4,5,6} All these factors require thoughtful policy interventions that will allow Ugandan children the opportunity to thrive and lead healthy and productive lives.

Mental Health: Prevention and Early Intervention

When screened in Ugandan primary care clinics, 12 to 29% of children present mental health symptoms.^{7,8} More specifically, in a study of depression amongst adolescents in secondary schools in Uganda, Nalugya (2004)⁸ found that 21% of youth presented depression symptoms. The prevalence of anxiety disorders has been found to be as high as 26.6%, with rates higher in females (29.7%) than in males (23.1%).⁹ Adolescent suicidality in Uganda has also been high.¹⁰

Mental health challenges are associated with increased risk for poverty due to factors such as increased health expenses, compromised productivity, mental health stigma, and loss of employment.

Mental health challenges that emerge during childhood and adolescence may compromise healthy transition to adulthood and increase economic and social costs for families, governments, and society in general.^{11,12} For instance, childhood disruptive behavior disorders, if not addressed, are associated with adverse outcomes, including academic problems (e.g., school dropout), social impairment, a higher incidence of chronic physical problems, unemployment and legal problems, substance abuse and violence as adults.^{13,14} Moreover, studies show that a substantial proportion of mental health challenges in adults originate during childhood and adolescence.¹⁵ Hence, addressing mental health challenges in early developmental stages has been set as a priority for the global child health agenda.¹⁵



RECOMMENDATION #1

Include language in the current Mental Health Bill that prioritizes children and adolescents.

The Mental Health Bill provides an opportunity to address the needs of children in Uganda. Specific language that identifies child and adolescent mental health as one of the key priority areas would highlight the critical need of all Ugandan children to have access to quality mental health care.

RECOMMENDATION #2

Early detection can ensure long-term health and socioeconomic benefits.

Policies must explicitly address strengthening capacity for addressing child and adolescent mental health care needs in non-stigmatizing settings, including families, schools, and primary health care clinics.¹⁶ Government is best positioned to successfully embed early detection and care within existing child serving systems through the passage of the Mental Health Bill.

RECOMMENDATION #3

The Mental Health Bill should be renamed The Mental Health Care Bill.

The pending legislation should be renamed the Mental Health Care Bill to reflect Parliament's commitment to caring for the needs of our children, families, and communities. This will also allow to make the language less stigmatizing for people impacted by mental health challenges.

Endnotes

1. UNICEF (2015). State of the world's children 2015 country statistical tables: Uganda statistics. *UNICEF*, Retrieved from: http://www.unicef.org/infobycountry/uganda_statistics.html
2. Population Reference Bureau (2009). World population data sheet. *Population Reference Bureau*. Retrieved from: http://www.prb.org/pdf09/09wpds_eng.pdf.
3. World Health Organization (2009). Country profile of environmental burden of disease: Uganda. *World Health Organization*. Retrieved from: http://www.who.int/quantifying_ehimpacts/national/countryprofile/uganda.pdf?ua=1.
4. Koenig, M.A., Lutalo, T., Zhao, F., Nalugoda, F., Wabwire-Mangen, F., Kiwanuka, N., Wagman, J., Serwadda, D., Wawer, M., & Gray, R. (2003). Domestic violence in rural Uganda: Evidence from a community-based study. *Bulletin of the World Health Organization* 81(1), 53-60.
5. Naker, D. (2005). Violence against children: The voices of Uganda children and adults. *Kampala: Raising Voices and Save the Children in Uganda*, 1-116.
6. Ovuga, E., Boardman, J., & Wasserman, D. (2005). The prevalence of depression in two districts of Uganda. *Social Psychiatry and Psychiatric Epidemiology*, 40(6), 439-445. doi: 10.1007/s00127-005-0915-0
7. Giel, R., & Harding, T. W. (1976). Psychiatric priorities in developing countries. *The British Journal of Psychiatry*, 128(6), 513-522. doi: 10.1192/bjp.128.6.513
8. Nalugya J. (2004). Depression amongst secondary school adolescents in Mukono district, Uganda (Doctoral Dissertation). Kampala: Makerere University.
9. Abbo, C., Kinyanda, E., Kizza, R. B., Levin, J., Ndyabangi, S., & Stein, D. J. (2013) Prevalence, comorbidity and predictors of anxiety disorders in children and adolescents in rural north-eastern Uganda. *Child and Adolescent Psychiatry Ment Health*. 7(1), 21. doi: 10.1186/1753-2000-7-21. PMID: 23841918.
10. Ashenafi, Y., Kebede, D., Desta, M., & Alem, A. (2001). Prevalence of mental and behavioural disorders in Ethiopia. *East African Medical Journal*, 78(6), 308-311.
11. Leibson, C. L., Katusic, S. K., Barbaresi, W. J., Ransom, J., & O'Brien, P.C. (2001) Use and costs of medical care for children and adolescents with and without Attention-Deficit/Hyperactivity Disorder. *Journal of the American Medical Association (JAMA)*, 285(1), 60-66. doi:10.1001/jama.285.1.60
12. Scott, S. (2002) Parent training programmes. In M. Rutter, & E. Taylor (Eds.), *Child and Adolescent Psychiatry* (4th edn) (pp. 949-967). Oxford: Blackwell Publishing.
13. Belfer, M. L. (2008). Child and adolescent mental disorders: The magnitude of the problem across the globe. *Journal of Child Psychology and Psychiatry*, 49(3), 226-236. doi: 10.1111/j.1469-7610.2007.01855.x
14. Lendingham, J. E. (1999) Children and adolescents with oppositional defiant disorder and conduct disorder in the community: Experiences at school and with peer. In H. E. Quay, A. E. Hogan (Eds.), *Handbook of Disruptive Behavior Disorders* (pp. 353-370). New York: Plenum Press.
15. Kessler, R. C., Angermeyer, M., Anthony, J. C., De Graaf, R. O. N., Demyttenaere, K., Gasquet, I., ... & Kawakami, N. (2007). Lifetime prevalence and age-of-onset distributions of mental disorders in the World Health Organization's World Mental Health Survey Initiative. *World Psychiatry*, 6(3), 168-76. PMID: PMC2174588.
16. Kieling, C., Baker-Henningham, H., Belfer, M., Conti, G., Ertem, I., Omigbodun, O., Rohde, L. A., Srinath, S., Ulkuer, N., & Rahman, A. (2011). Child and adolescent mental health worldwide: Evidence for action. *The Lancet*, 378(9801), 1515-1525. doi: 10.1016/S0140-6736(11)60827-1

Acknowledgment: Part of this policy work is supported by the National Institute of Mental Health (NIMH) under Award Number U19 MH110001 (MPIs: Fred Ssewamala, PhD; Mary McKay, PhD; Kimberly Hoagwood, PhD). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Mental Health, the National Institutes of Health or Washington University in St. Louis.

Child and Adolescent Mental Health Care in Uganda

This report was developed by ICHAD (International Center for Child Health and Development), SMART Africa Center (Strengthening Mental health And Research Training in Sub-Saharan Africa), the Clark-Fox Policy Institute at the Brown School at Washington University in St. Louis and ChildFund Uganda.



In Uganda, children make up about half (56%) of the total population, and they often present with multiple physical, mental health, and educational challenges.^{1,2} Large numbers of Ugandan children live in communities with high rates of chronic poverty (38%), domestic violence (30%), physical violence toward children (80%), depression (33 to 39%), malaria (70 to 80%), and HIV or AIDS (6%).^{3,4,5,6} All these factors require thoughtful policy interventions that will allow Ugandan children the opportunity to thrive and lead healthy and productive lives.

Workforce Training

Uganda, like many other developing countries, has scarce mental health workforce resources (e.g., psychiatrists, psychologists, social workers and nurses). Moreover, very few of these professionals are trained specifically in child and adolescent mental health care.⁷ Given the limited number of mental health care professionals in Uganda (and most SSA countries), it is not feasible to rely solely on such professionals to deliver mental health prevention and care services. Although primary health care professionals can provide the bulk of care, mental health professionals, namely psychiatrists, nurses and experts in psychosocial health, are also needed to provide adequate services to those children and adolescents who require intensive intervention. Additionally, the mental health care workforce plays a key role in delivering training, support and supervision to non-

specialists. Without these mental health care professionals, Uganda will not have enough human resources to meet their populations' mental health treatment requirements, including children and adolescents. This is even more urgent given that Ugandan population primarily comprises of children and adolescents (56%).

RECOMMENDATION #1

Primary health care workers must be trained in child and adolescent mental health.

Research evaluating the effectiveness of mental health training for various groups, including general practitioners and nurses found that the trainings were especially successful in improving diagnostic skills and sensitivity of trainees as well as their attitudes towards mental health challenges.⁸

RECOMMENDATION #2

Additional training is needed for mental health care workers.

Given the unique needs of children and adolescents with mental health challenges, supplemental training programs, (e.g., certificate programs, advanced degree programs) for the primary mental health care workforce can add to the pool of individuals trained in child and adolescent mental health.⁹ Training in the delivery of brief, low-burden, evidence-based interventions adapted to the Ugandan context has been found effective, feasible, and acceptable.⁸

RECOMMENDATION #3

Support and train lay workers and peers that already exist in health and education systems to implement evidence-based mental health interventions.

Task-shifting (also referred to as task sharing), endorsed by the World Health Organization, is a cost-efficient and feasible model that involves redistributing tasks from professionally trained workers to those with less training and fewer qualifications.⁹ Research has demonstrated that mental health interventions provided by local lay counselors with little to no previous mental health training or experience have demonstrated positive findings in the area of mental health, health, and overall psychosocial outcomes.⁹ Community health workers, village health teams, expert clients -and others who may fall in that category- who work directly with children and families at Health Center 1 and LC 1 levels could constitute this workforce in the Ugandan context.

Endnotes

1. UNICEF (2015). State of the world's children 2015 country statistical tables: Uganda statistics. *UNICEF*, Retrieved from: http://www.unicef.org/infobycountry/uganda_statistics.html
2. Population Reference Bureau (2009). World population data sheet. *Population Reference Bureau*. Retrieved from: http://www.prb.org/pdf09/09wpds_eng.pdf.
3. World Health Organization (2009). Country profile of environmental burden of disease: Uganda. *World Health Organization*. Retrieved from: http://www.who.int/quantifying_ehimpacts/national/countryprofile/uganda.pdf?ua=1.
4. Koenig, M.A., Lutalo, T., Zhao, F., Nalugoda, F., Wabwire-Mangen, F., Kiwanuka, N., Wagman, J., Serwadda, D., Wawer, M., & Gray, R. (2003). Domestic violence in rural Uganda: Evidence from a community-based study. *Bulletin of the World Health Organization* 81(1), 53-60.
5. Naker, D. (2005). Violence against children: The voices of Uganda children and adults. *Kampala: Raising Voices and Save the Children in Uganda*, 1-116.
6. Ovuga, E., Boardman, J., & Wasserman, D. (2005). The prevalence of depression in two districts of Uganda. *Social Psychiatry and Psychiatric Epidemiology*, 40(6), 439-445. doi: 10.1007/s00127-005-0915-0
7. Bass, J., Neugebauer, R., Clougherty, K. F., Verdelli, H., Wickramaratne, P., Ndogoni, L., ... & Bolton, P. (2006). Group interpersonal psychotherapy for depression in rural Uganda: 6-month outcomes: randomised controlled trial. *The British Journal of Psychiatry*, 188(6), 567-573.
8. Ayano, G., Assefa, D., Haile, K., Chaka, A., Haile, K., Solomon, M., ... Jemal, K. (2017). Mental health training for primary health care workers and implication for success of integration of mental health into primary care: evaluation of effect on knowledge, attitude and practices (KAP). *International Journal of Mental Health Systems*, 11, 63. <http://doi.org/10.1186/s13033-017-0169-8>
9. Patel, V., Weiss, H. A., Chowdhary, N., Naik, S., Pednekar, S., Chatterjee, S., ... & Simon, G. (2010). Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *The Lancet*, 376(9758), 2086-2095.

Acknowledgment: Part of this policy work is supported by the National Institute of Mental Health (NIMH) under Award Number U19 MH110001 (MPIs: Fred Ssewamala, PhD; Mary McKay, PhD; Kimberly Hoagwood, PhD). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Mental Health, the National Institutes of Health or Washington University in St. Louis.

Child and Adolescent Mental Health Care in Uganda

This report was developed by ICHAD (International Center for Child Health and Development), SMART Africa Center (Strengthening Mental health And Research Training in Sub-Saharan Africa), the Clark-Fox Policy Institute at the Brown School at Washington University in St. Louis and ChildFund Uganda.



In Uganda, children make up about half (56%) of the total population, and they often present with multiple physical, mental health, and educational challenges.^{1,2} Large numbers of Ugandan children live in communities with high rates of chronic poverty (38%), domestic violence (30%), physical violence toward children (80%), depression (33 to 39%), malaria (70 to 80%), and HIV or AIDS (6%).^{3,4,5,6} All these factors require thoughtful policy interventions that will allow Ugandan children the opportunity to thrive and lead healthy and productive lives.

Asset-Based Economic Development Aimed at Addressing Poverty

According to the World Bank, Uganda reduced monetary poverty from 31.1% in 2006 to 19.7% in 2013.⁷ However, the proportion of the Ugandan children and adolescents living below the national poverty has reached 38% for children under five and 55% for children ages six to 17. Of those children, 18% live in extreme poverty and face serious challenges to healthy development and transition to adulthood.¹ Poverty during childhood also increases the risk for intergenerational poverty.⁸

Research has pointed to the bidirectional relationship between poverty and mental health, with poverty increasing the probability of mental health challenges through heightened stress, social exclusion, decreased social capital, malnutrition, and increased obstetric risks, violence, and trauma.⁹ In turn, mental health challenges are associated with increased risk for poverty due to

factors such as increased health expenses, compromised productivity, mental health stigma, and loss of employment/unemployment.¹⁰

Children and adolescents in communities assaulted by persistent poverty and disease are more likely to suffer from a range of chronic mental health challenges, which in turn, undermine their ability to become functional members of the society. Poverty has also been found to negatively impact children's brain/cognitive development.¹¹

Studies have pointed to the bidirectional relationship between poverty and mental health, with poverty increasing the probability of mental illness.

RECOMMENDATION #1

Asset-based interventions can play a critical role in reducing risks associated with mental health challenges.

Economic empowerment interventions, including family and child savings accounts, need to be offered to poverty-impacted adolescents and families in Uganda. Research has demonstrated that poor families in low-resource communities can effectively benefit economically from these interventions. Additionally, the interventions can minimize risk taking behaviors and address mental health stressors among children and adolescents.^{12, 13, 14, 15, 16, 17}

RECOMMENDATION #2

Providing financial literacy training to children and adolescents can promote saving behavior.

Integrating financial management trainings into school curriculum can contribute to teaching and promoting saving behavior among children and adolescents, which in turn, may reduce poverty and its associated risks.^{12, 13, 14, 15, 16, 17}

Endnotes

1. UNICEF (2015). State of the world's children 2015 country statistical tables: Uganda statistics. *UNICEF*, Retrieved from: http://www.unicef.org/infobycountry/uganda_statistics.html
2. Population Reference Bureau (2009). World population data sheet. *Population Reference Bureau*. Retrieved from: http://www.prb.org/pdf09/09wpds_eng.pdf.
3. World Health Organization (2009). Country profile of environmental burden of disease: Uganda. *World Health Organization*. Retrieved from: http://www.who.int/quantifying_ehimpacts/national/countryprofile/uganda.pdf?ua=1.
4. Koenig, M.A., Lutalo, T., Zhao, F., Nalugoda, F., Wabwire-Mangen, F., Kiwanuka, N., Wagman, J., Serwadda, D., Wawer, M., & Gray, R. (2003). Domestic violence in rural Uganda: Evidence from a community-based study. *Bulletin of the World Health Organization* 81(1), 53-60.
5. Naker, D. (2005). Violence against children: The voices of Uganda children and adults. *Kampala: Raising Voices and Save the Children in Uganda*, 1-116.
6. Ovuga, E., Boardman, J., & Wasserman, D. (2005). The prevalence of depression in two districts of Uganda. *Social Psychiatry and Psychiatric Epidemiology*, 40(6), 439-445. doi: 10.1007/s00127-005-0915-0
7. The World Bank (2016). Uganda poverty assessment 2016: Fact sheet. *The World Bank*. Retrieved from: <http://www.worldbank.org/en/country/uganda/brief/uganda-poverty-assessment-2016-fact-sheet>
8. Bird, K. (2007). The intergenerational transmission of poverty: An overview. Chronic Poverty Research Centre Working Paper No. 99. doi: 10.2139/ssrn.1629262
9. Flisher, A. J., Lund, C., Funk, M., Banda, M., Bhana, A., Doku, V., ... & Petersen, I. (2007). Mental health policy development and implementation in four African countries. *Journal of Health Psychology*, 12(3), 505-516.
10. Saraceno, B., Levav, I., Kohn, R. (2005). The public mental health significance of research on socio-economic factors in schizophrenia and major depression. *World Psychiatry*, 4(3), 181-85.
11. Luby, J., Belden, A., Botteron, K., Marrus, N., Harms, M. P., Babb, C., ... & Barch, D. (2013). The Effects of poverty on childhood brain development: The mediating effect of caregiving and stressful life events. *Journal of the American Medical Association Pediatrics*, 167(12), 1135-1142. doi: 10.1001/jamapediatrics.2013.3139
12. Ssewamala, F. M., & Ismayilova, L. (2009). Integrating children's savings accounts in the care and support of orphaned adolescents in rural Uganda. *Social Service Review*, 83(3), 453-472. PMID: PMC2863345
13. Ssewamala, F. M., Han, C-K., Neilands, T. B., Ismayilova, L., & Sperber, E. (2010). Effect of economic assets on sexual risk-taking intentions among orphaned adolescents in Uganda. *American Journal of Public Health*, 100(3), 483-488. PMID: PMC2820050
14. Ssewamala, F.M., Nabunya, P., N.Miraim, M., Ilic, V. & Nattabi, J. (2014) Integrating a mentorship component in programming for care and support of AIDS-Orphaned and Vulnerable Children: Lessons from the Suubi and Bridges Programs in Sub-Saharan Africa. *Global Social Welfare*, 1, 9-24. doi: 1007/s40609-014-0008-7
15. Nabunya, P., Ssewamala, F. M., & Ilic, V. (2014). Family economic strengthening and parenting stress among caregivers of AIDS-orphaned children: Results from a cluster randomized clinical trial in Uganda. *Children and Youth Service Review*, 44, 417-421. PMID: PMC4133737
16. Ssewamala, F. M., Nabunya, P., Ilic, V., Mukasa, M., & Ddamulira, C. (2015). Relationship between family economic resources, psychosocial well-being, and educational preferences of AIDS-orphaned children in southern Uganda: Baseline findings. *Global Social Welfare*, 2(2), 75-86. doi: 10.1007/s40609-015-0027-z
17. Curley, J., Ssewamala, F. M., Nabunya, P., Ilic, V., & Han, C-K. (2014). Child Development Accounts (CDAs): An asset-building strategy to empower girls in Uganda. *International Social Work*, 57(2), 1-14.

Acknowledgment: Part of this policy work is supported by the National Institute of Mental Health (NIMH) under Award Number U19 MH110001 (MPIs: Fred Ssewamala, PhD; Mary McKay, PhD; Kimberly Hoagwood, PhD). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Mental Health, the National Institutes of Health or Washington University in St. Louis.